Crohn's Disease Ulcerative Colitis (drugs A-R)

(Cimzia[®], Humira®)



There owned There owned There of the services Specialty Services There of the services The service owned The service own

Sex: □ Female Male SSN:	Patient Information Prescriber + Shipping Information										
Language: Wt: Dkg Dibs Ht: Dem Din Address: Address:	Patient name: DOB:				Prescrib	Prescriber name:					
Address:											
Address:	Language: Wt: Dkg Dlbs Ht:Dcm Din Address:										
Phone:			Apt/Suite	e:	City:	St	ate:	Zip:			
Caregiver name: Relation:	Apt/Suite: City:					:					
Local pharmacy: Phone: Insurance plan: Plan ID: Insurance plan: Plan ID: If shipping to prescriber: First Fill Always Never (Corbn's Disease of the insurance card(s). (Initical Information (Places fax all partitical and fab information) (Corbn's Disease of the Small Intensities) (K50.9 (Crohn's Disease of Both Intensities) (K50.9 (Crohn's Disease of Both Intensities) (K50.9 (Crohn's Disease of Both Intensities) (K51.9 (Ulcerative Rectosigmoiditis) (K51.3 (Ulcerative Rectosigmoiditis) (K51.3 (Ulcerative Colitis) (Coronitations) (Linget 10 ong subcut at weeks 0, 2 and 4 (Autive Colitis) (Inject 40 ong subcut					Phone: Alternate:						
Local pharmacy: Phone: Insurance plan: Plan ID: Insurance plan: Plan ID: If shipping to prescriber: First Fill Always Never (Corbn's Disease of the insurance card(s). (Initical Information (Places fax all partitical and fab information) (Corbn's Disease of the Small Intensities) (K50.9 (Crohn's Disease of Both Intensities) (K50.9 (Crohn's Disease of Both Intensities) (K50.9 (Crohn's Disease of Both Intensities) (K51.9 (Ulcerative Rectosigmoiditis) (K51.3 (Ulcerative Rectosigmoiditis) (K51.3 (Ulcerative Colitis) (Coronitations) (Linget 10 ong subcut at weeks 0, 2 and 4 (Autive Colitis) (Inject 40 ong subcut											
Insurance plan: Plan 10: If shipping to prescriber: First Fill Always Never Please fax a copy of front and back of the insurance card(s). Cincleal Information (Please Raill pertinent clinical and fabs information) Crohn's Disease of the Small Intenstine)K50.0 (Crohn's Disease of the Large Intenstine) K50.0 (Crohn's Disease of Both Intenstines)K50.0 (Crohn's Disease) of the Large Intenstine) K50.0 (Crohn's Disease of Both Intenstines)K50.0 (Crohn's Disease) of the Large Intenstine) K50.0 (Crohn's Disease of Both Intenstines)K50.0 (Crohn's Disease) of the Large Intenstine) K51.3 (Ulcerative Rectosigmoidins) K51.3 (Ulcerative Rectosigmoidins) K51.3 (Ulcerative Rectosigmoidins) K51.3 (Ulcerative Rectosigmoidins) K51.3 (Ulcerative Rectosigmoidins) Cronorbiant Medications: Concomitant Medications: Concomitant Medications: Concomitant Medications: Concomitant Medications: Cincala® (certolizumab) Aubits Inject 400 mg subcut at weeks 0, 2 and 4 F x 200 mg/mL PFS Inject 400 mg subcut on day 1, then 80 mg on day 15 Inject 400 mg subcut on day 1, then 80 mg on day 15 Inject 400 mg subcut day 1, then 80 mg on day 15 PFS Inject 400 mg subcut on day 20 and every other week thereafter 2 x 40 mg/0.8mL PFS Inject 160 mg subcut day 1, then 80 mg on day 15											
Please fax a copy of front and back of the insurance card(s). Clinical Information (Please fax all portinent clinical and lab information) Crohn's Disease: K80.0 (Crohn's Disease of the Small Intenstine) Crohn's Disease: K80.0 (Crohn's Disease of the Small Intenstine) Crohn's Disease: K80.0 (Crohn's Disease of the Small Intenstine) Crohn's Disease: K80.0 (Crohn's Disease of the Small Intenstine) Crohn's Disease: K80.0 (Crohn's Disease) Constraints K51.5 (Leff Sided Colitis) K51.5 (Leff Sided Colitis) K51.8 (Other Ulcerative Rectosigmoiditis) Diagnosis Date: TB Test: Yes< No Neg. Test Date:											
Clinical information (Clease fax all pertinent clinical and lab information) K50.1 (Crohn's Disease of the Small Intensities) K50.1 (Crohn's Disease of the Small Intensities) K50.1 (Crohn's Disease, unspecified) Utcerative Colitis: K51.0 (Ucerative Pancolitis) K51.3 (Ucerative Rectosigmoiditis) K51.5 (Left Sided Colitis) K51.8 (Other Ucerative Colitis) K51.9 (Ucerative Colitis) Diagnosis Date: TB Test: Yes No Prior Therapy Yes No Reason for Discontinuation of Therapy Approximate End Date Concomitant Medications: Reason for Discontinuation of Therapy Approximate End Date	Please fax a copy of front and back of the insurance card(s).										
Crohn's Disease of the Small Intensitienel K50.1 (Crohn's Disease of the Large Intensitienel) K50.4 (Crohn's Disease of Both Intensitienel) K50.9 (Crohn's Disease, unspecified) Ulcerative Colitis: K51.0 (Ulcerative Pancolitis) K51.3 (Ulcerative Rectosigmoiditis) K51.5 (Left Sided Colitis) K51.8 (Other Ulcerative Colitis) K51.3 (Ulcerative Rectosigmoiditis) Diagnosis Date: TB Test: Yes No Prior Therapy Yes No Reason for Discontinuation of Therapy Approximate Start Date Prior Therapy Yes No Reason for Discontinuation of Therapy Approximate Start Date Comorbidities: Comorbidities: Comorbidities: Presscription Comorbidities: Other: PPS Vials 0 Inject 400 mg subcut at weeks 0, 2 and 4 6 x 200 mg/mL PPS Vials 0 Inject 400 mg subcut on day 1, then 80 mg on day 15 0 for to 40 kg) PFS 0 Inject 400 mg subcut on day 29 and every other 2 x 40 mg/0.8mL Pens 0 Prestinter® Inject 40 mg subcut day 1, then 80 mg on day 15 3 x 40 mg/0.8mL PFS 0 Inject 40 mg subcut on day 29 and every other 2											
Ulcerative Colitis: K51.0 (Ulcerative Pancolitis) K51.3 (Ulcerative Rectosigmoiditis) Other: K51.5 (Left Sided Colitis) K51.8 (Other Ulcerative Colitis) K51.9 (Ulcerative Colitis) Other: TB Test: Yes No K51.3 (Ulcerative Colitis) K51.9 (Ulcerative Colitis) Other: TB Test: Yes No K51.3 (Ulcerative Colitis) K51.9 (Ulcerative Colitis) Originosis Date: TB Test: Yes No Neg. Test Date: Approximate Start Date Prior Therapy Yes No Reason for Discontinuation of Therapy Approximate Start Date Approximate End Date Comorbidities: Concontiant Medications: Allergies: No Pens 0 Cortorizana® Inject 400 mg subcut at weeks 0, 2 and 4 6 x 200 mg/mL PFS 0 Inject 160 mg subcut on day 1, then 80 mg on day 15 6 x 40 mg/0.8mL Pens 0 Aduits Inject 400 mg subcut on day 29 and every other week thereafter 2 x 40 mg/0.8mL Pens 0 Inject 40 mg subcut on day 29 and every other week thereafter 3 x 40 mg/0.8mL Pens 0 Inject 20 mg subcut on day 29 and every other week thereafter (17 to <40 kg)	Crohn's Disease: K50.0 (Crohn's Disease of the Small Intenstine)										
Abst.5 (Left Sided Colitis) K51.8 (Other Ulcerative Colitis) K51.9 (Ulcerative Colitis) Other: TB Test: Yes No Neg. Test Date: Prior Therapy I Yes No Reason for Discontinuation of Therapy Prior Therapy Prior Therapy Yes NKDA NKDA Ocnornitant Medications: Concontinant Medications: Inject 400 mg subcut at weeks 0, 2 and 4 I higet 160 mg subcut on day 1, then 80 mg on day 15 I higet 160 mg subcut on day 29 and every other I higet 160 mg subcut day 1, then 80 mg on day 15 (17 to <40 kg) I higet 160 mg subcut day 1, then 80 mg on day 15 (17 to <40 kg) I higet 160 mg subcut day 1, then 80 mg on day 15 (2 x 40 mg/0.8mL Pers I higet 160 mg subcut day 1, then 80 mg on day 15 (17 to <40 kg) I higet 160 mg subcut day 1, then 80 mg on day 15 (2 x 40 mg/0.8mL Pers I higet 160 mg subcut day 1, then 80 mg on day 15 (2 x 40 mg/0.8mL Pers I higet 160 mg subcut day 1, then 80 mg on day 15 (2 40 kg) I higet 160 mg subcut day 1, then 80 mg on day 15 (2 x 40 mg/0.8mL Pers I higet 160 mg subcut on day 29 and every other Week thereafter I higet 20 mg subcut on day 29 and every other Week thereafter (17 to <40 kg) I higet 20 mg subcut on day 29 and every other Week thereafter (17 to <40 kg) I higet 20 mg subcut on day 29 and every other Week thereafter (240 kg) I higet 20 mg subcut on day 29 and every other Week thereafter (240 kg) I higet 20 mg subcut on day 29 and every other Week thereafter (240 kg) I higet 20 mg subcut on day 29 and every other </td <td colspan="10"></td>											
Other:											
Diagnosis Date: TB Test: Yes No Neg. Test Date: Approximate Start Date Approximate End Date Prior Therapy Yes No Reason for Discontinuation of Therapy Approximate Start Date Approximate End Date Comorbidities:											
Prior Therapy Yes No Reason for Discontinuation of Therapy Approximate Start Date Approximate End Date											
Comorbidities:						ie		Start Data	Approvimat	o End Dato	
Concomitant Medications: Allergies: NKDA □ Other: Prescription □ Inject 400 mg subcut at weeks 0, 2 and 4 □ 6 x 200 mg/mL □ PFS □ Cimzia [®] □ Inject 400 mg subcut every 4 weeks □ 2 x 200 mg/mL □ PFS □ Linject 160 mg subcut on day 1, then 80 mg on day 15 □ 1nject 400 mg subcut on day 29 and every other week thereafter □ 2 x 40 mg/0.8mL □ Pens □ □ Adults □ Inject 400 mg subcut day 1, then 40 mg on day 15 □ 1nject 80 mg subcut day 1, then 40 mg on day 15 □ 1 x 40 mg/0.8mL □ PFS □ □ Humira [®] □ Inject 80 mg subcut day 1, then 40 mg on day 15 □ 3 x 40 mg/0.8mL □ Pens □ □ Inject 80 mg subcut on day 29 and every other week thereafter □ 1 sto 40 kg) □ 1nject 160 mg subcut on day 29 and every other □ 3 x 40 mg/0.8mL □ Pens 0 □ Rediatincs ≥ 6 years □ Inject 20 mg subcut on day 29 and every other week thereafter (17 to <40 kg)			Reason for Disco		егару	Approximate Start Date A			Аррголіта	e chu Dale	
Concomitant Medications: Allergies: NKDA □ Other: Prescription □ Inject 400 mg subcut at weeks 0, 2 and 4 □ 6 x 200 mg/mL □ PFS □ Cimzia [®] □ Inject 400 mg subcut every 4 weeks □ 2 x 200 mg/mL □ PFS □ Linject 160 mg subcut on day 1, then 80 mg on day 15 □ 1nject 400 mg subcut on day 29 and every other week thereafter □ 2 x 40 mg/0.8mL □ Pens □ □ Adults □ Inject 400 mg subcut day 1, then 40 mg on day 15 □ 1nject 80 mg subcut day 1, then 40 mg on day 15 □ 1 x 40 mg/0.8mL □ PFS □ □ Humira [®] □ Inject 80 mg subcut day 1, then 40 mg on day 15 □ 3 x 40 mg/0.8mL □ Pens □ □ Inject 80 mg subcut on day 29 and every other week thereafter □ 1 sto 40 kg) □ 1nject 160 mg subcut on day 29 and every other □ 3 x 40 mg/0.8mL □ Pens 0 □ Rediatincs ≥ 6 years □ Inject 20 mg subcut on day 29 and every other week thereafter (17 to <40 kg)				<u>-</u>							
Concomitant Medications: Allergies: NKDA □ Other: Prescription □ Inject 400 mg subcut at weeks 0, 2 and 4 □ 6 x 200 mg/mL □ PFS □ Cimzia [®] □ Inject 400 mg subcut every 4 weeks □ 2 x 200 mg/mL □ PFS □ Linject 160 mg subcut on day 1, then 80 mg on day 15 □ 1nject 400 mg subcut on day 29 and every other week thereafter □ 2 x 40 mg/0.8mL □ Pens □ □ Adults □ Inject 400 mg subcut day 1, then 40 mg on day 15 □ 1nject 80 mg subcut day 1, then 40 mg on day 15 □ 1 x 40 mg/0.8mL □ PFS □ □ Humira [®] □ Inject 80 mg subcut day 1, then 40 mg on day 15 □ 3 x 40 mg/0.8mL □ Pens □ □ Inject 80 mg subcut on day 29 and every other week thereafter □ 1 sto 40 kg) □ 1nject 160 mg subcut on day 29 and every other □ 3 x 40 mg/0.8mL □ Pens 0 □ Rediatincs ≥ 6 years □ Inject 20 mg subcut on day 29 and every other week thereafter (17 to <40 kg)											
Allergies: D NKDA □ Other:	Comorbidities:										
Prescription Cimzia® Inject 400 mg subcut at weeks 0, 2 and 4 6 x 200 mg/mL Vials 0 Inject 400 mg subcut every 4 weeks 2 x 200 mg/mL Vials 0 Inject 400 mg subcut on day 1, then 80 mg on day 15 0 Vials 0 Adults Inject 40 mg subcut on day 1, then 80 mg on day 15 0 PFS 0 Inject 40 mg subcut on day 29 and every other week thereafter 10 a x 40 mg/0.8mL Pens 0 Inject 80 mg subcut day 1, then 40 mg on day 15 (17 to <40 kg)	Concomitant Medication	ons:									
Image: Cimzia® (certolizumab) Inject 400 mg subcut at weeks 0, 2 and 4 Image: 6 x 200 mg/mL Image: Viais 0 Image: Inject 400 mg subcut every 4 weeks Image: 2 x 200 mg/mL Image: Viais Image: Viais 0 Image: Inject 400 mg subcut on day 1, then 80 mg on day 15 Image: Inject 400 mg subcut on day 29 and every other week thereafter Image: 2 x 40 mg/0.8mL Image: Pers event evet	Allergies: 🛛 NKDA 🗆	Other:									
□ Cimzia® (certolizumab) □ Inject 400 mg subcut at weeks 0, 2 and 4 □ 6 x 200 mg/mL □ Vials 0 □ Inject 400 mg subcut every 4 weeks □ 2 x 200 mg/mL □ PFS □ Vials □ PFS □ Vials 0 □ Humira® (adalimumab) □ Inject 400 mg subcut on day 1, then 80 mg on day 15 □ 6 x 40 mg/0.8mL □ Pens □ FFS 0 □ Adults □ Inject 400 mg subcut on day 29 and every other week thereafter □ 2 x 40 mg/0.8mL □ Pens □ PFS 0 □ Humira® (adalimumab) □ Inject 80 mg subcut day 1, then 40 mg on day 15 (17 to <40 kg)	Prescription							1			
Inject 400 mg subcut every 4 weeks □ 2 x 200 mg/mL □ Vials Inject 160 mg subcut on day 1, then 80 mg on day 15 □ Pens 0 (adalimumab) □ Inject 40 mg subcut on day 29 and every other week thereafter □ 2 x 40 mg/0.8mL □ Pens 0 □ Humira® □ Inject 80 mg subcut day 1, then 40 mg on day 15 □ 2 x 40 mg/0.8mL □ Pens 0 □ Humira® □ Inject 80 mg subcut day 1, then 40 mg on day 15 □ 3 x 40 mg/0.8mL □ Pens 0 □ Humira® □ Inject 160 mg subcut day 1, then 80 mg on day 15 □ 3 x 40 mg/0.8mL □ Pens 0 □ Inject 20 mg subcut on day 29 and every other week thereafter (17 to <40 kg)	□ Cimzia [®]	□ Inject 400 mg subcut at weeks 0, 2 and 4				□ 6 x 200 mg/mL				0	
□ Humira® day 15 □ 6 x 40 mg/0.8mL □ Pens 0 □ Adults □ Inject 40 mg subcut on day 29 and every other □ 2 x 40 mg/0.8mL □ PFS □ □ Adults □ Inject 40 mg subcut day 1, then 40 mg on day 15 □ 2 x 40 mg/0.8mL □ PFS □ □ Humira® □ Inject 80 mg subcut day 1, then 40 mg on day 15 □ 3 x 40 mg/0.8mL □ Pens □ □ Humira® □ Inject 160 mg subcut day 1, then 80 mg on day □ 3 x 40 mg/0.8mL □ Pens 0 □ Adults □ Inject 160 mg subcut day 1, then 80 mg on day □ 3 x 40 mg/0.8mL □ Pens 0 □ floct 160 mg subcut on day 29 and every other □ 2 x 20 mg/0.8mL □ Pens 0 □ Inject 20 mg subcut on day 29 and every other □ 2 x 20 mg/0.4mL PFS	(certolizumab)	Inject 400 mg subcut every 4 weeks				□ 2 x 200 mg/mL		_			
□ Humira® day 15 □ 6 x 40 mg/0.8mL □ Pens 0 □ Adults □ Inject 40 mg subcut on day 29 and every other □ 2 x 40 mg/0.8mL □ PFS □ □ Adults □ Inject 40 mg subcut day 1, then 40 mg on day 15 □ 2 x 40 mg/0.8mL □ PFS □ □ Humira® □ Inject 80 mg subcut day 1, then 40 mg on day 15 □ 3 x 40 mg/0.8mL □ Pens □ □ Humira® □ Inject 160 mg subcut day 1, then 80 mg on day □ 3 x 40 mg/0.8mL □ Pens 0 □ Adults □ Inject 160 mg subcut day 1, then 80 mg on day □ 3 x 40 mg/0.8mL □ Pens 0 □ floct 160 mg subcut on day 29 and every other □ 2 x 20 mg/0.8mL □ Pens 0 □ Inject 20 mg subcut on day 29 and every other □ 2 x 20 mg/0.4mL PFS		□ Iniect 160	ma subcut on day ?	1. then 80 ma o	n						
(adalimumab) □ <	□ Humira [®]	day 15				□ 6 x 40 mg/0.8mL				0	
Aduits □ Inject 40 mg subcut on day 29 and every other week thereafter □ 2 x 40 mg/0.8mL □ Pens □ □ Humira® □ Inject 80 mg subcut day 1, then 40 mg on day 15 (17 to <40 kg)											
□ Inject 10 mg outbout on day 20 and overy other □ 2 × 40 mg/0.8mL □ PFS □ Humira® □ Inject 80 mg subcut day 1, then 40 mg on day 15 (17 to <40 kg)	. ,					-		Pens			
□ Inject 80 mg subcut day 1, then 40 mg on day 15 (17 to <40 kg)	, louino					□ 2 x 40 mg/0.8mL					
Image: (17 to <40 kg)					v 15						
□ Humira® (adalimumab) □ Inject 160 mg subcut day 1, then 80 mg on day 15 (≥40 kg) □ PFS 0 □ Inject 20 mg subcut on day 29 and every other week thereafter (17 to <40 kg)		(17 to <40 kg)			-						
□ Humira® 15 (≥40 kg) □										0	
(adalimumab) □	□ Humira [®]						ng/0.8mL				
week thereafter (17 to <40 kg)	(adalimumab)		·								
□ Inject 40 mg subcut on day 29 and every other week thereafter (≥40 kg) □ 2 x 40 mg/0.8mL □ Pens § Simponi [®] and Stelara [®] are available on the Crohn's Disease/Ulcerative Colitis Enrollment Form S-Z § Injection Training Provided by: □ Physician's Office □ Pharmacy □ Other: Per state-specific law, prescriptions will be dispensed as generic, if applicable, unless notated otherwise:	Pediatrics ≥ 6 years					□ 2 x 20 mg/0.4mL		PFS			
week thereafter (≥40 kg) □ PFS § Simponi [®] and Stelara [®] are available on the Crohn's Disease/Ulcerative Colitis Enrollment Form S-Z § Injection Training Provided by: □ Physician's Office □ Pharmacy □ Other: Per state-specific law, prescriptions will be dispensed as generic, if applicable, unless notated otherwise:		□ Inject 40 mg subcut on day 29 and every other						Pens			
Injection Training Provided by: Physician's Office Pharmacy Other: Per state-specific law, prescriptions will be dispensed as generic, if applicable, unless notated otherwise:				,		_		PFS			
Injection Training Provided by: Physician's Office Pharmacy Other: Per state-specific law, prescriptions will be dispensed as generic, if applicable, unless notated otherwise:	§ Simponi [®] and Stelara [®] are available on the Crohn's Disease/Ulcerative Colitis Enrollment Form S-Z §										
Per state-specific law, prescriptions will be dispensed as generic, if applicable, unless notated otherwise:	Injection Training Provided by: Dephysician's Office Department Other:										
			-	-			tated otherwise	e:			
Stanip Signature not anowed, physician Signature required.											
Prescriber's Signature: Date:											
I authorize Thrifty White Specialty Pharmacy and its representatives to act as an agent to initiate and execute the insurance prior authorization process for this prescription and any future fills of the same prescription for the patient listed above. I understand that I can revoke this designation at any time by providing written notice to Thrifty White Specialty Pharmacy.											

Confidentiality Statement: This message is intended only for the individual or entity to which it is addressed. It may contain information which may be proprietary and confidential. It may also contain privileged, confidential information which is exempt from disclosure under applicable laws, including the Health Insurance Portability and Accountability Act (HIPAA). If you are not the intended recipient, please note that you are strictly prohibited from disseminating or distributing this information (other than to the intended recipient) copying this information. If you received this communication in error, please notify the sender immediately by calling 855-611-3399 or by emailing specialty@thriftywhite.com to obtain instructions as to the proper destruction of the transmitted material. Thank you.